

Probabilistic interpretation of compositeness relation for resonances

Zhi-Hui Guo^{1,2} and J. A. Oller³

¹ Department of Physics, Hebei Normal University, Shijiazhuang 050024, People's Republic of China

² State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, CAS, Beijing 100190, People's Republic of China

³ Departamento de Física, Universidad de Murcia, E-30071 Murcia, Spain

Bound, antibound and resonance states are associated to poles in the on-shell partial wave amplitudes. We show here that from the residues of the pole a rank 1 projection operator associated with any of these states can be extracted, in terms of which a sum rule related to the composition of the state can be derived. Although typically it involves complex coefficients for the compositeness and elementariness, except for the bound state case, we demonstrate that one can formulate a meaningful compositeness relation with only positive coefficients for resonances whose associated Laurent series in the variable s converges in a region of the physical axis around Res_P , with s_P the pole position of the resonance. It is also shown that this result can be considered as an analytical extrapolation in s_P of the clear narrow resonance case. We exemplify this formalism to study the two-body components of several resonances of interest.

I. INTRODUCTION

Resonance, as it arises in quantum-mechanical problems, is a common phenomenon in several branches of physics, e.g. in particle, nuclear, atomic and condensed matter physics. Here we make use of S matrix theory, but the same principles hold in other disciplines as well. An interesting question in spectroscopy is to understand the nature of resonances, so that one can see whether they can fit within the standard model of physics or they have another origin. Even in the former case, the situation requires further clarification in many instances in order to understand their nature in terms of the appropriate degrees of freedom.

During the past decades there has been an increasing growth of evidences that hadron resonances do not always fit within the standard picture for mesons ($q\bar{q}$) and baryons (qqq) in quantum chromodynamics. Well known examples are the early puzzles of the $\Lambda(1405)$ and related resonances, the lightest scalar mesons, and the examples of new bottomonium, charmonium and charm-strange mesons with unexpected properties. In this respect, one also has the exciting new discovery of the $P_c(4450)$ [1] that would require five valence quarks. All these examples clearly show that the interest on hadron resonances and their nature is actually reinvigorated.

A tool to understand the composition of a state is through the field renormalization constant Z , first formulated for shallow bound states [2], and that represents the amount of extra components beyond the explicit degrees of freedom considered ($1 - Z$ is then called the compositeness). Its generalization to resonances lying nearby a threshold is discussed in Refs. [3, 4]. We focus here on another approach based on the analytical continuation of the compositeness relation to the pole position [5–7]. This approach allows a probabilistic interpretation for two-body composition of bound states, but for the case of resonances its straightforward application drives to complex numbers which have prevented such interpretation. We show here how this difficulty can be overcome by employing appropriate transformations driving to a new unitary S matrix that shares the same resonant behavior but gives rise to compositeness relation involving only positive numbers.

II. SUM RULE

In the following we consider on-shell two-body scattering of n channels, and assume rotational and time reversal symmetry, so that the S and T matrices are symmetric [8]. The T matrix can be written in general form as [9]

$$T(s) = [\mathcal{K}(s)^{-1} + G(s)]^{-1}, \quad (1)$$

where s is the usual Mandelstam variable and $G(s)$ is a diagonal matrix with matrix elements $\delta_{ij}G(s)_i$. The $G(s)_i$ are the unitarity scalar-loop functions that encode the two-body unitarity requirement, $\text{Im}T^{-1}(s)_{ij} = -\delta_{ij}\theta(s-s_i)\rho(s)_i$, as well as its analytical properties. These functions can be expressed as [10]

$$G(s)_i = a(s_0)_i - \frac{s-s_0}{\pi} \int_{s_i}^{\infty} ds' \frac{\rho(s')_i}{(s'-s)(s'-s_0)}, \quad (2)$$

and s_i is the threshold for the i_{th} channel and we order the states such that $s_i < s_{i+1}$. In the previous expressions $\rho(s)_i$ is the relativistic invariant phase space factor for channel i , $\rho(s)_i = q(s)_i/8\pi\sqrt{s}$, with $q(s)_i$ the center-of-mass three-momentum. We denote $\rho(s)$ as the diagonal matrix, with matrix elements $\delta_{ij}\rho(s)_i$. In addition, the matrix $\mathcal{K}(s)$ in Eq. (1) includes all other contributions not arising from the two-body unitarity, such as the contact interactions and the crossed-channel effects. In our notation the relation between the S and T matrices is

$$S(s) = I + i(2\rho(s))^{\frac{1}{2}}T(s)(2\rho(s))^{\frac{1}{2}}, \quad (3)$$

with I the identity matrix. Equation (3) can also be applied (through analytical continuation) even if s is below the threshold for some of the channels because its restriction to the subspace of open channels gives the correct S matrix.

Next we assume that the T matrix has a pole with a given set of quantum numbers in the appropriate Riemann sheet (RS) at s_P . This pole is called a bound state if its lies in the physical RS, $s_P \in \mathbb{R}$ and $s_P < s_1$; an antibound state if $s_P \in \mathbb{R}$, $s_P < s_1$ but it lies in the unphysical RS adjacent to the physical one for $s \gtrsim s_1$; and a resonance state in any other case (in the

following we denote all of them as pole states). The residues of the T matrix at s_P are given by

$$\lim_{s \rightarrow s_P} (s - s_P)T(s) = -\gamma\gamma^T, \quad (4)$$

where γ is an n row vector, $\gamma^T = (\gamma_1, \dots, \gamma_n)$, with γ_i typically complex numbers. Now, by employing the algebraic result that $dB(s)^{-1}/ds = -B^{-1}(s)dB(s)/dsB^{-1}(s)$, with $B(s)$ an invertible matrix, we take the derivative with respect to s of both sides of Eq. (1) at $s = s_P$. From the double-pole term at s_P it follows that

$$\gamma\gamma^T = -\gamma\gamma^T \left[\frac{dG(s_P)}{ds} - \mathcal{K}^{-1} \frac{d\mathcal{K}(s_P)}{ds} \mathcal{K}^{-1} \right] \gamma\gamma^T. \quad (5)$$

Let us stress that the derivative of $G(s)_i$ is independent of the subtraction constant $a(s_0)_i$ and subtraction point s_0 . It corresponds to a convergent three-point one-loop function with unit vertices [11]. Next, we multiply Eq. (5) to the left by $\gamma^\dagger$ and to the right by γ^* , and as $\gamma^\dagger \gamma \neq 0$ it can be simplified. Furthermore, in the last term on the right-hand side (rhs) of Eq. (5) we employ that $\gamma = -\mathcal{K}G\gamma$, as follows by rewriting Eq. (1) as $T = \mathcal{K} - \mathcal{K}GT$ and taking the limit $s \rightarrow s_P$. Then, one obtains the sum rule (SR)

$$1 = -\gamma^T \frac{dG(s_P)}{ds} \gamma + \gamma^T G(s_P) \frac{d\mathcal{K}(s_P)}{ds} G(s_P) \gamma \quad (6)$$

$$= \sum_{i,j=1}^n \left(-\delta_{ij} \gamma_i^2 \frac{dG(s_P)_i}{ds} + \gamma_i G(s_P)_i \frac{d\mathcal{K}(s_P)_{ij}}{ds} G(s_P)_j \gamma_j \right).$$

This SR was already derived in the literature [5–7]. The partial compositeness coefficients from Eq. (6) are given by $X_i = -\gamma_i^2 dG(s_P)_i/ds$, their sum is the total compositeness, $X = \sum_{i=1}^n X_i$, and the elementariness (the analytically continued field renormalization constant at the pole), $Z = 1 - X$, corresponds to the sum over the last term on the rhs of Eq. (6). For bound states all these coefficients are positive and their connection with the different components in the properly normalized bound state is understood [2, 3, 7].¹ However, for the rest of the pole states Eq. (6) does not allow a straightforward statistical quantum mechanical interpretation since the compositeness and elementariness coefficients become usually complex.

A. Projection operator of rank 1

Around the pole position we can write for the partial-wave projected S matrix the Laurent expansion

$$S(s) = \frac{\mathcal{R}}{s - s_P} + S_0(s), \quad (7)$$

where $\mathcal{R}$ is the matrix of residues and $S_0(s)$ comprises the nonresonant terms. From Eqs. (3) and (4) one has the following relation between the residues of the S and T matrices,

$$\mathcal{R} = -2i\rho(s_P)^{\frac{1}{2}} \gamma\gamma^T \rho(s_P)^{\frac{1}{2}}. \quad (8)$$

Let us show next that the multiplicative structure of the residues implies the existence of a projection operator of rank 1. For that we rewrite the matrix $\mathcal{R}$ in an explicitly symmetric form as

$$\mathcal{R} = i\lambda \mathcal{O} \mathcal{A} \mathcal{O}^T, \quad (9)$$

with λ a real normalization constant. Combining Eqs. (8) and (9) leads to

$$\mathcal{A} = -\frac{1}{\lambda} \omega \omega^T, \quad (10)$$

where

$$\omega = \mathcal{O}^{-1} \tilde{\gamma}, \quad (11)$$

$$\tilde{\gamma} = (2\rho(s_P))^{\frac{1}{2}} \gamma.$$

Next we require that $\mathcal{A}$ is a projection operator, so that $\mathcal{A}^\dagger = \mathcal{A}$ and $\mathcal{A}^2 = \mathcal{A}$. The former condition implies that the vector ω must be real or purely imaginary (these are n constraints on the $2n^2$ free parameters present in $\mathcal{O}$), and we take it real in the following (this can always be done because $\mathcal{A}$ depends quadratically on ω). Next, the latter condition fixes λ to

$$\lambda = -\omega^T \omega < 0. \quad (12)$$

This result also implies that $\mathcal{A}$ is of rank 1 because its trace is

$$\text{Tr} \mathcal{A} = -\frac{1}{\lambda} \text{Tr} \omega \omega^T = 1. \quad (13)$$

Notice that given a projection $\mathcal{A}$ with the above properties and L a real orthogonal matrix then $L\mathcal{A}L^T$ shares the same properties as $\mathcal{A}$.

A deeper understanding of the SR in Eq. (6) can be reached by rewriting $\gamma\gamma^T$ in Eq. (5) in terms of $\mathcal{A}$ according to Eq. (10). After simplifying common factors one has

$$\mathcal{A} = \frac{\lambda}{2} \mathcal{A} \mathcal{O}^T \rho(s_P)^{-\frac{1}{2}} \left[\frac{dG(s_P)}{ds} - G(s_P) \frac{d\mathcal{K}(s_P)}{ds} G(s_P) \right] \times \rho(s_P)^{-\frac{1}{2}} \mathcal{O} \mathcal{A}. \quad (14)$$

Next, we take the trace on both sides of this equation, employ on the rhs the cyclic property of the trace and then Eq. (6) results. Although the connection of this SR with the proper normalization of bound states was already derived [2, 3], now we understand that it is also equal to 1 for resonances and antibound states because of the proper normalization (to 1) of the pole state associated with the rank 1 projection operator $\mathcal{A}$.

¹ There is also another proposal in the literature for higher partial waves [6] that consists of using a redefined T matrix in order to get the SR. However, this is physically equivalent, and leads to the same results, both on the physical axis and for the values of X_i , by properly defining the accompanying matrices $\mathcal{K}(s)$ and $G(s)$ [12].

B. Open and closed channels

Let us follow the convention to write a hat when denoting submatrices in the reduced space with m open channels ($s_m < s < s_{m+1}$) and, when indicated, the subscript m corresponds to the number of them. Since again $\hat{S}_m(s)$ is a unitarity matrix it has its associated T matrix, $\hat{T}_m(s)$, which is the restriction of the original one in the subspace of open channels, and its residues are denoted by $\hat{\gamma}$. The matrix $\hat{T}_m(s)$ can be written analogously to Eq. (1) in terms of an effective $\hat{\mathcal{K}}_m(s)$, which can be calculated from the original $\mathcal{K}(s)$ by integrating out heavier channels [7]. We can also derive the subsequent SR in the reduced subspace, which reads

$$1 = \hat{\gamma}^T \left[-\frac{d\hat{G}_m(s_P)}{ds} + \hat{G}_m(s_P) \frac{d\hat{\mathcal{K}}_m(s_P)}{ds} \hat{G}_m(s_P) \right] \hat{\gamma}. \quad (15)$$

This new SR can be related to the original one, expressed for n channels, by introducing an $m \times n$ matrix $\mathcal{H}$ such that

$$\mathcal{H} = \begin{pmatrix} \hat{I}_m & 0 \end{pmatrix}, \quad (16)$$

$$\hat{T}_m(s) = -\frac{i}{2} \hat{\rho}(s)^{-\frac{1}{2}} \left[\mathcal{H} S(s) \mathcal{H}^T - \hat{I}_m \right] \hat{\rho}(s)^{-\frac{1}{2}}, \quad (17)$$

where $\hat{G}_m(s_P) = \mathcal{H} G(s)$ and $\hat{\gamma} = \mathcal{H} \gamma$, with $\hat{I}_m$ the $m \times m$ identity matrix. As done above, the SR results by taking the derivative of Eq. (17) at s_P ,

$$\begin{aligned} & \hat{\gamma} \hat{\gamma}^T \left[\hat{G}_m(s_P) \frac{d\hat{\mathcal{K}}_m(s_P)}{ds} \hat{G}_m(s_P) - \frac{d\hat{G}_m(s_P)}{ds} \right] \hat{\gamma} \hat{\gamma}^T \\ &= \hat{\gamma} \gamma^T \left[G(s_P) \frac{d\mathcal{K}(s_P)}{ds} G(s_P) - \frac{dG(s_P)}{ds} \right] \gamma \gamma^T, \end{aligned} \quad (18)$$

and multiplying next both sides to the left by $\hat{\gamma}^\dagger$ and to the right by $\hat{\gamma}^*$. As the product $\hat{\gamma}^\dagger \hat{\gamma} \neq 0$ the equality between the SRs of Eqs. (6) and (15) follows. Note that since the X_i , $i = 1, \dots, m$, are the same in both schemes, from the equality between these SRs one obtains the relation

$$\hat{Z} = \hat{\gamma}^T \hat{G}_m \frac{d\hat{\mathcal{K}}_m(s_P)}{ds} \hat{G}_m \hat{\gamma} = Z + \sum_{i=m+1}^n X_i. \quad (19)$$

Thus, the elementariness $\hat{Z}$ in the reduced space contains also the contribution from two-body heavier channels [7]. This result is derived here in a fully general manner without referring to any specific interaction.

III. TRANSFORMED SUM RULES

We point out that the SR in Eq. (15) is not unique. Given an $m \times m$ unitary matrix $\hat{U}_m$ we define the transformation

$$\hat{S}_u(s) = \hat{U}_m \hat{S}_m(s) \hat{U}_m^T. \quad (20)$$

This S matrix is symmetric and also unitary for $s_m < s < s_{m+1}$, so that one can identify a transformed T matrix, $\hat{T}_u(s)$ (given by the rhs of Eq. (17) replacing $\mathcal{H} \rightarrow \hat{U}_m$ and $S \rightarrow \hat{S}_m$), which could be expressed as in Eq. (1) in terms of a new $\hat{\mathcal{K}}_u(s)$. The matrix of residues for $\hat{S}_u(s)$ is $\hat{\mathcal{R}}_u = \hat{U}_m \hat{\mathcal{R}} \hat{U}_m^T$ or, in other words, the matrix $\hat{\mathcal{O}}$, cf. Eq. (9), transforms to $\hat{\mathcal{O}}_u = \hat{U}_m \hat{\mathcal{O}} \hat{U}_m^T$, while the projection operator $\hat{\mathcal{A}}_m$ does as $\hat{L} \hat{\mathcal{A}}_m \hat{L}^T$ (with $\hat{L}$ the aforementioned real orthogonal matrix). In terms of the residues of $\hat{T}_u(s)$, $\hat{\gamma}_u = \hat{\rho}(s_P)^{-\frac{1}{2}} \hat{U}_m \hat{\rho}(s_P)^{\frac{1}{2}} \hat{\gamma}$, and $\hat{\mathcal{K}}_u(s)$ a new SR attached to $\hat{\mathcal{A}}_m$ can be derived

$$1 = -\hat{\gamma}_u^T \frac{d\hat{G}_m(s_P)}{ds} \hat{\gamma}_u + \hat{\gamma}_u^T \hat{G}_m \frac{d\hat{\mathcal{K}}_u(s_P)}{ds} \hat{G}_m \hat{\gamma}_u. \quad (21)$$

A. New completeness relation

In the subsequent all our results are based on the hypothesis that the pole s_P lies in an unphysical RS adjacent to the physical RS in an interval $s_m < s < s_{m+1}$, such that $s_m < \text{Res}_P < s_{m+1}$. We define the resonance mass squared, M_R^2 , as $M_R^2 = \text{Res}_P$. The important point is that, as a result, the convergence region of the Laurent series in Eq. (7) includes a region of the real axis around $s = M_R^2$ (since this point is closer to s_P than any of the nearest thresholds s_m or s_{m+1}).² We denote this working assumption as **I**. In particular it excludes from our considerations the case of anti-bound states or resonances that lie in the RS mentioned but with $M_R^2 < s_m$ or $M_R^2 > s_{m+1}$ (e.g. this is the case of the $a_0(980)$ in Refs. [10, 13]. These types of resonant signals are an admixture between a pole and an enhanced cusp effect by the pole itself [14]). Condition **I** is certainly satisfied by a narrow resonance. But we should stress that **I** goes far beyond the narrow resonance case (the latter is discussed in detail below in Sec. III B).

Based on the condition **I** we can further inquire about the matrix $\mathcal{O}$ in Eq. (9). Since we are going to make use of the unitarity of the S matrix around $s = M_R^2$, we consider its restriction $\hat{S}_m(s)$ in the subset of the m open channels. Correspondingly, the analysis undertaken above from Eq. (9) until concluding the existence of the rank 1 projection operator is restricted to this subset as well. Performing a transformation of the type of Eq. (20) we would have

$$\hat{S}_u(s) = \hat{U}_m \hat{S}_m(s) \hat{U}_m^T = \frac{i\lambda \hat{\mathcal{A}}_m}{s - s_P} + \hat{U}_m \hat{S}_0(s) \hat{U}_m^T, \quad (22)$$

where $i\lambda \hat{\mathcal{A}}_m = \hat{U}_m \hat{\mathcal{R}}_m \hat{U}_m^T$. As both matrices $\hat{\mathcal{R}}_m$ and $\hat{\mathcal{A}}_m$ are constant we consider only constant $\hat{U}_m$. This is why

² For resonances lying far enough in the complex plane it is not excluded that the Laurent series does not actually converge in this energy region because of the closeness of some singularity due to crossed channel dynamics. This depends on the particular dynamics affecting every independent process and should be studied case by case.

we disregarded any energy dependence for the transformation matrix in Eq. (20).

Now, the point is that because condition **I** holds then we can move from the real axis onto the pole position (due to the convergence of the Laurent series for s around M_R^2) and identify the matrix $\mathcal{O}$ with $\hat{U}_m^{-1}$. Since it is a unitary transformation the real vector $\hat{\omega}$, given by $\hat{\omega} = \hat{U}_m(2\rho(s_P))^{\frac{1}{2}}\hat{\gamma}$, must have a modulus squared equal to

$$\hat{\omega}^T \hat{\omega} = \sum_{i=1}^m |\tilde{\gamma}_i|^2. \quad (23)$$

This constraint cannot be determined from the analysis undergone above, which concluded with the existence of the projection operator $\mathcal{A}$, because there we only attended to the factorizing structure of the residues, while here we assume that **I** holds. Indeed, given a real vector $\hat{\omega}$ with such modulus squared any other one transformed by an arbitrary real orthogonal matrix $\hat{L}_m$, $\hat{\omega}' = \hat{L}_m \hat{\omega}$, is equally valid mathematically. Then, we can always take the unitary transformation $\hat{U}_m$ such that the components of $\hat{\omega}$ are

$$\omega_i = |\tilde{\gamma}_i|, \quad i = 1, \dots, m. \quad (24)$$

This is accomplished e.g. by the diagonal unitary matrix $\hat{U}_m = \text{diag}(\frac{|\tilde{\gamma}_1|}{\tilde{\gamma}_1}, \dots, \frac{|\tilde{\gamma}_m|}{\tilde{\gamma}_m})$, with the understanding that if $\tilde{\gamma}_i = 0$ one takes 1 instead of $\frac{|\tilde{\gamma}_i|}{\tilde{\gamma}_i}$ in $\hat{U}_m$. The main motivation to finally fix the action of $\hat{L}_m$ in this way (as a result $\hat{\mathcal{A}}_m$ is invariant under the transformation of Eq. (20)) is to avoid mixing between different channels in the resulting matrix of residues for the resonance, leaving intact the original strength (modulus) of the resonance coupling to each channel. Hence we can say that the resonance signal around the pole is preserved by the transformation (from the narrow resonance case it is well known that the phases of the residues are due to the background [8, 12, 15], cf. Sec. IIIB). In this way the physical picture for the pole term in Eq. (7) as corresponding to the resonance exchange between channels driving by the original residues is preserved.

For the T matrix corresponding to Eqs. (22) and (24) the residues are $|\gamma_i|\eta_i$, with η_i a pure phase factor given by $|\rho(s_P)_i|/\rho(s_P)_i$. However, the new coefficients entering in the resulting SR, analogous to Eq. (21), are not still positive definite because of the complex numbers $dG(s_P)_i/ds$ and η_i . We then allow by an extra unitary transformation of the kind in Eq. (20) given by a constant diagonal matrix $\hat{U}'_m = \text{diag}(e^{i\phi_1}, \dots, e^{i\phi_m})$, so that no mixing between resonance couplings to different channels arises. The residues of the new T matrix are $|\gamma_i|e^{i\phi_i}\eta_i$, and the phase factors $e^{i\phi_i}$ are fixed by requiring that the resulting partial compositeness coefficients, X_i^R , $i = 1, \dots, m$, are positive. Hence,

$$X_i^R = -e^{2i\phi_i}\eta_i^2|\gamma_i|^2 \frac{dG(s_P)_i}{ds} = |\gamma_i|^2 \left| \frac{dG(s_P)_i}{ds} \right|. \quad (25)$$

In terms of them one has the total compositeness, $X^R = \sum_{i=1}^n X_i^R$, and elementariness $Z^R = 1 - X^R$. One should

stress that these coefficients are entirely determined from the properties of the pole of the resonance (s_P and γ_i) and no further ambiguity remains in the appropriate SR for a resonant state fulfilling **I**, as the proper transformation of Eq. (20) has been fixed. This also implies that $X^R \leq 1$ should hold since the right SR must be unique. At this point it is worth stressing that if the T matrix were known in a region (no matter how small) of the physical s axis around M_R^2 , where the Laurent series Eq. (7) converges, one-variable complex analysis guarantees that the different contributions in this series (in particular s_P and γ_i) could be determined unambiguous and model independently.

Let us stress that we are relying on unitarity and analyticity, cf. Eq. (22), and then our results could be applied to a T matrix with correct analytical properties in the complex s plane, in particular along the unitarity cut. In this respect it is especially important to keep the whole energy dependence for $G(s)_i$ as dictated by Eq. (2).³ In particular the restriction $X^R \leq 1$ should be satisfied, similarly as s_P , γ and S_0 are linked to satisfy unitarity on the real axis wherever the Laurent series Eq. (7) converges. It is clear that there is a close connection between narrow resonances (in which the two-body open channels are trapped for a relatively long time) and bound states, so that the compositeness relation for both cases should meet. Nevertheless, for a bound state both γ_i^2 and $-dG(s_P)_i/ds$ are positive and one does not need to take the absolute value in Eq. (25) [5–7].

B. Narrow resonance case

We now assume that the resonance dominates the energy dependence of the expansion in Eq. (7) on the real axis around $s = M_R^2$, such that we can take in good approximation $S_0(s)$ as a constant matrix (this is typically the narrow resonance assumption). To simplify the writing we also drop the hat on top of the different matrices, so that one should understand in the following that the analysis is restricted to the set of m open channels.

Implementing in the unitarity relation $S(s)S(s)^\dagger = 1$ the expansion in Eq. (7) we have the equation

$$(s - s_P)(s - s_P^*)S_0S_0^\dagger + (s - s_P)S_0R^\dagger + (s - s_P^*)RS_0^\dagger + RR^\dagger = (s - s_P)(s - s_P^*). \quad (26)$$

By identifying the coefficients of the different powers of s we have the following set of equations

$$S_0S_0^\dagger = I, \quad (27)$$

$$S_0R^\dagger + RS_0^\dagger = 0, \quad (28)$$

$$s_P S_0 R^\dagger + s_P^* R S_0^\dagger - R R^\dagger = 0. \quad (29)$$

³ For its nonrelativistic reduction see Ref. [16].

The first of them implies that S_0 is a unitary matrix. One can find a family of solutions of these equations by writing

$$S_0 = \mathcal{O} \mathcal{O}^T, \quad (30)$$

$$R = i\lambda \mathcal{O} \mathcal{A} \mathcal{O}^T, \quad \lambda \in \mathbb{R}, \quad (31)$$

with $\mathcal{O}$ a unitary matrix, so that Eq. (27) is fulfilled. Although S_0 is symmetric the matrix $\mathcal{O}$ in Eq. (30) is not necessarily symmetric.⁴ We have from Eqs. (28) and (29) that the matrix $\mathcal{A}$ satisfies, respectively,

$$i\lambda \mathcal{A} - i\lambda \mathcal{A}^\dagger = 0, \quad (32)$$

$$-2\text{Im } s_P \mathcal{A} + \lambda \mathcal{A} \mathcal{A}^\dagger = 0. \quad (33)$$

From the first equation it is clear that

$$\mathcal{A}^\dagger = \mathcal{A}, \quad (34)$$

while we can always take the normalization constant λ such that additionally Eq. (33) is fulfilled with

$$\mathcal{A}^2 = \mathcal{A}, \quad (35)$$

$$\lambda = 2\text{Im } s_P = -2M_R \Gamma_R. \quad (36)$$

As a result of Eqs. (34) and (35) it follows that $\mathcal{A}$ is a projection operator. It is also a symmetric matrix, cf. Eq. (31), and then its matrix elements must be real because it is a Hermitian matrix, cf. Eq. (34). Additionally, $\mathcal{A}$ is a rank 1 projector operator by direct analogy between Eqs. (9) and (31). We can prove this conclusion explicitly here for the narrow resonance case by noticing that

$$\text{Tr} \mathcal{A} = -\frac{1}{\lambda} \text{Tr} |\tilde{\gamma}| |\tilde{\gamma}|^T = \frac{\sum_{i=1}^m |\gamma_i|^2 |\rho(s_R)_i|}{M_R \Gamma_R} = 1. \quad (37)$$

The equality to 1 of this sum stems from the replacement of $\rho(s_R) \rightarrow \rho(M_R^2)$ in Eq. (37), since $\Gamma_R \ll M_R$, and the standard formula for the total decay width of a narrow resonance [16],

$$\Gamma_R = \sum_{i=1}^m \frac{|\gamma_i|^2 q(M_R^2)_i}{8\pi M_R^2}. \quad (38)$$

Then, for the narrow resonance case we can write the S matrix, Eq. (7), around the resonance mass as

$$S(s) = \mathcal{O} \left(I + \frac{i\lambda \mathcal{A}}{s - s_P} \right) \mathcal{O}^T, \quad (39)$$

with $\mathcal{A}$ a rank 1 projection operator and λ given by Eq. (36). For a given projection $\mathcal{A}$ one can identify a resonance S matrix, $S_R(s)$, given by the matrix between brackets in the previous equation, namely,

$$S_R(s) = I + \frac{i\lambda \mathcal{A}}{s - s_P}. \quad (40)$$

However, this identification is not unique because $\mathcal{A}$ can be changed by the action of a real orthogonal matrix, as already discussed after Eq. (13), $\mathcal{A} \rightarrow L\mathcal{A}$ and $\mathcal{O} \rightarrow \mathcal{O}L^T$. Since $\mathcal{O}$ is unitary Eq. (23) necessarily holds and L is finally fixed by the requirement of Eq. (24), $\omega_i = |\tilde{\gamma}_i|$. In this way, the resonance projection operator is determined and from this analysis it is clear that the phases of the original couplings γ_i are due to the nonresonant terms through the unitary matrix $\mathcal{O}$, cf. Eqs. (30) and (39).⁵

Nevertheless, one has to make still one more transformation of the type of Eq. (22) to guarantee that the resulting compositeness coefficients X_i , $i = 1, \dots, m$, are real. This extra transformation is accomplished by a diagonal unitary matrix, analogously as done above in Eq. (25), and is needed mainly because $dG(s_P)_i/ds$, $i = 1, \dots, m$ are complex, cf. footnote 5. In the transformed S matrix, $S_u(s)$, the unitary matrix $\mathcal{O}$ transforms as $\mathcal{O} \rightarrow \mathcal{U}\mathcal{O}$, while the resonance S matrix, $S_R(s)$, remains untouched.

In summary, our derivations in Secs. II A and III can be considered as the generalization of the narrow resonance case by extrapolating analytically in the resonance pole position s_P from the real axis at M_R^2 (that corresponds to the limit of a narrow resonance $\Gamma_R/M_R \rightarrow 0$) up to s_P , along a contour that runs parallel to the imaginary s axis. However, this extrapolation is only possible while **I** holds. Among other facts notice that $dG(s)_m/ds$ has as branch-point singularity at s_m , and let us recall that $s_m < M_R^2 < s_{m+1}$. If condition **I** is not met one enters in a new qualitative physical picture that manifests by the fact that the Laurent series of Eq. (7) does not converge for any real s .

C. General $\mathcal{O}$ matrix

When **I** does not hold we cannot conclude that $\hat{\mathcal{O}}$, cf. Eqs. (9) and (22), corresponds to a unitary matrix $\hat{\mathcal{U}}_m^{-1}$. A further insight into this problem can be gained by writing a left polar decomposition for $\hat{\mathcal{O}}^{-1}$,

$$\hat{\mathcal{O}}^{-1} = Q\mathcal{V}, \quad (41)$$

where $\mathcal{V}$ is a unitary matrix and Q is positive definite Hermitian matrix.⁶ The matrix Q can be further diagonalized as

$$Q = \mathcal{Z} Q_d \mathcal{Z}^\dagger \quad (42)$$

where $Q_d = \text{diag}(d_1, \dots, d_n)$ ($d_i > 0$) and $\mathcal{Z}$ is a unitary matrix. In this form Eq. (41) and the vector ω in Eq. (11) read

$$\hat{\mathcal{O}}^{-1} = \mathcal{Z} Q_d \mathcal{Z}^\dagger \mathcal{V}, \quad (43)$$

$$\omega = \mathcal{Z} Q_d \mathcal{Z}^\dagger \mathcal{V} \tilde{\gamma}. \quad (44)$$

⁵ One has to take into account that $\rho(s_P)_i$, $i = 1, \dots, m$ are real in the narrow resonance limit $\Gamma_R/M_R \rightarrow 0$.

⁶ Let us recall that $\hat{\mathcal{O}}$ has inverse and this is why Q is positive definite and not simply positive-semidefinite Hermitian matrix.

⁴ A particular solution would be $\mathcal{O} = S_0^{1/2}$.

Name of the states	Pole: $\sqrt{s_P}$ [MeV]	$X_{\pi\pi}^R$	X_{KK}^R	$X_{\eta\eta}^R$	$X_{\eta\eta'}^R$	X^R	Z^R
$f_0(500)$ [18]	$442_{-4}^{+4} - i246_{-5}^{+7}$	$0.40_{-0.02}^{+0.02}$	...	...	...	$0.40_{-0.02}^{+0.02}$	$0.60_{-0.02}^{+0.02}$
$f_0(980)$ [18]	$978_{-11}^{+17} - i29_{-11}^{+9}$	$0.02_{-0.01}^{+0.01}$	$0.65_{-0.26}^{+0.27}$	...	...	$0.67_{-0.27}^{+0.28}$	$0.33_{-0.27}^{+0.28}$
$f_0(1710)$ [14]	$1690_{-20}^{+20} - i110_{-20}^{+20}$	$0.00_{-0.00}^{+0.00}$	$0.03_{-0.01}^{+0.01}$	$0.02_{-0.01}^{+0.01}$	$0.20_{-0.07}^{+0.07}$	$0.25_{-0.10}^{+0.10}$	$0.75_{-0.10}^{+0.10}$
$\rho(770)$ [18]	$760_{-5}^{+7} - i71_{-5}^{+4}$	$0.08_{-0.01}^{+0.01}$	...	...	...	$0.08_{-0.01}^{+0.01}$	$0.92_{-0.01}^{+0.01}$
$K_0^*(800)$ [18]	$643_{-30}^{+75} - i303_{-75}^{+25}$	$0.94_{-0.52}^{+0.39}$				$0.94_{-0.52}^{+0.39}$	$0.06_{-0.52}^{+0.39}$
$K^*(892)$ [18]	$892_{-7}^{+5} - i25_{-2}^{+2}$	$0.05_{-0.01}^{+0.01}$				$0.05_{-0.01}^{+0.01}$	$0.95_{-0.01}^{+0.01}$
$a_0(1450)$ [18]	$1459_{-95}^{+70} - i174_{-100}^{+110}$	$0.09_{-0.07}^{+0.03}$	$0.02_{-0.02}^{+0.12}$	$0.12_{-0.09}^{+0.22}$		$0.23_{-0.18}^{+0.37}$	$0.77_{-0.18}^{+0.37}$
$a_1(1260)$ [19]	$1260 - i250$	0.46				0.46	0.54
Hyperon with $I = 0$		$X_{\pi\Sigma}^R$	X_{KN}^R			X^R	Z^R
$\Lambda(1405)$ broad [20]	$1388_{-9}^{+9} - i114_{-25}^{+24}$	$0.73_{-0.10}^{+0.15}$	...			$0.73_{-0.10}^{+0.15}$	$0.27_{-0.10}^{+0.15}$
$\Lambda(1405)$ narrow [20]	$1421_{-2}^{+3} - i19_{-5}^{+8}$	$0.18_{-0.08}^{+0.13}$	$0.82_{-0.17}^{+0.36}$			$1.00_{-0.25}^{+0.49}$	$0.00_{-0.25}^{+0.49}$
Hyperon with $I = 1$		$X_{\pi\Lambda}^R$	$X_{\pi\Sigma}^R$	X_{KN}^R		X^R	Z^R
Ref. [20]	$1376_{-3}^{+3} - i33_{-5}^{+5}$	$0.04_{-0.00}^{+0.00}$	$0.0_{-0.0}^{+0.0}$	...		$0.04_{-0.00}^{+0.00}$	$0.96_{-0.00}^{+0.00}$
Ref. [20]	$1414_{-3}^{+2} - i12_{-2}^{+1}$	$0.03_{-0.00}^{+0.00}$	$0.01_{-0.00}^{+0.00}$	$0.13_{-0.04}^{+0.03}$		$0.17_{-0.04}^{+0.03}$	$0.83_{-0.04}^{+0.03}$
$D_{s0}^*(2317)$ [21]	2321_{-3}^{+6}	$0.56_{-0.03}^{+0.05}$	$0.12_{-0.01}^{+0.01}$	$0.02_{-0.01}^{+0.01}$		$0.70_{-0.05}^{+0.07}$	$0.30_{-0.05}^{+0.07}$
$Y(4260)$ [22, 23]	$4232.8 - i36.3$	$X_{J/\psi f_0(500)}^R$	$X_{J/\psi f_0(980)}^R$	$X_{Z_c(3900)\pi}^R$	$X_{\omega\chi_{c0}}^R$	X^R	Z^R
		0.00	0.02	0.02	0.17	0.21	0.79
$\Lambda_c(2595)$ [17]	$2592.25 - i1.3$	$X_{\Sigma^+\pi^0}^R$	$X_{\Sigma_c^+\pi^-}^R$	$X_{\Sigma_c^0\pi^+}^R$		X^R	Z^R
		$0.11_{-0.02}^{+0.02}$	...	...		$0.11_{-0.02}^{+0.02}$	$0.89_{-0.02}^{+0.02}$

TABLE I. Partial compositeness coefficients (X_i^R), total compositeness (X^R) and elementariness (Z^R) for a set of resonances that satisfy condition I.

For diagonal $\hat{\mathcal{O}}$, which is the one finally involved in Secs. III A and III B, and it is also necessarily the case for the important one-channel scattering, instead of Eq. (23) we have now

$$\hat{\omega}^T \hat{\omega} = \sum_{i=1}^m d_i |\tilde{\gamma}_i|^2. \quad (45)$$

There is no way to fix the positive constants d_i with just the information contained in the residue matrix $\mathcal{R}$ at the pole position s_P . However, when condition I holds we can relate directly the SR at the pole position with a transformed unitary S matrix $\hat{S}_u(s)$, cf. Eq. (22), and conclude that $\hat{\mathcal{O}}_m$ is unitary ($d_i = 1$, $i = 1, \dots, m$). In other words, $d_i \neq 1$ cannot be excluded when calculating compositeness if I is not met,⁷ and then $\hat{\mathcal{O}}$ does not correspond to the unitary-matrix-driven transformation of Eq. (20) from a resonance S matrix.

IV. APPLICATION TO SOME RESONANCES OF INTEREST

We proceed to calculate the values of compositeness and elementariness coefficients from Eq. (25) to examine the nature of a set of resonances that satisfy the condition I and whose pole position and residues are taken from literature. We should stress that our aim here is to exemplify our method for the calculation of X_i , cf. Eq. (25), for several resonances of interest but not to discern the goodness between possible different analyses/models for a given resonance nor being exhaustive in listing such analyses. We present values for the partial compositeness coefficients in Table I, with the channels involved appearing as subscripts. Several different types of resonances are considered, including light-flavor mesons, light-flavor baryons, and mesons and baryons with heavy flavors.

We briefly elaborate how the numbers in Table I are obtained. For a set of resonances both the pole positions and the relevant residues (central values and uncertainties), are explicitly given in the references taken, which enable us to straightforwardly calculate the compositeness X_i by using Eq. (25). The error bars given in Table I arise by propagating and adding in quadrature the errors from the pole position s_P and the residues γ_i^2 . We do not attempt here to make an estimation

⁷ As we have explicitly checked in many examples.

of possible sources of systematic errors in the theoretical approach followed by that reference. Namely, the set of resonances used, are: the light-flavor mesons $f_0(500)$, $f_0(980)$, $\rho(770)$, $K_0^*(800)$, $K^*(892)$, $a_0(1450)$ from Ref. [18], the $f_0(1710)$ state from Ref. [14], the isoscalar and isovector hyperons from Ref. [20], and the bound state $D_{s0}^*(2317)$ from Ref. [21]. For the $a_1(1260)$, we have taken into account the different normalization to properly relate the coupling strength in Ref. [19] with γ_i in this work.

Now we consider the heavy-flavor states. For the light-flavor resonances we observe that Z^R is clearly dominant for the $\rho(770)$ and $K^*(892)$ (as it should be for genuine $q\bar{q}$ resonances [18, 24]). To a lesser extent this is also the case for the $f_0(1710)$ (identified in Ref. [14] as mainly a glueball state) and the $a_0(1450)$ (which is a representative of the second nonet of scalar resonances with dominantly preexisting nature [10, 14, 18]). Regarding the lightest scalars we see that the two-body component is much more important, being overwhelmingly dominant for the $K_0^*(800)$ and the largest for the $f_0(980)$. Nevertheless, the compositeness for the $f_0(500)$ is more modest, although it raises with increasing pion mass, so that $X^R \simeq 0.6$ for $M_\pi \simeq 220$ MeV (while still satisfying condition I) [25]. For the axial-vector $a_1(1260)$ we obtain that X^R and Z^R are similar, which is compatible with Ref. [26] that finds an N_c pole trajectory at odds with a $q\bar{q}$ resonance. Regarding the light-flavor hyperon resonances, the $\Lambda(1405)$ is clearly a resonance mainly composed by the light-pseudoscalar mesons and the baryon octet, the lighter pole dominated by the $\pi\Sigma$ component and the heavier one by the $\bar{K}N$ component [27]. However, these two-body components are small for the two isovector-hyperon poles [9, 28].

Notice that we have calculated the compositeness of the $f_0(980)$ in $K\bar{K}$ and that of the $\Lambda(1405)$ in $\bar{K}N$, although I is not really fulfilled because these channels have a threshold larger than M_R^2 . Our calculation in these cases is based on the fact that the difference between M_R and this threshold is significantly smaller than the width. Thus, the transition through the threshold is rather smooth and then we can expect that $\omega_{m+1} \simeq |\tilde{\gamma}_{m+1}|$ in good approximation, cf. Eq. (24), since the transformation above s_{m+1} must be unitary. As a result we can apply Eq. (25) as an approximation for the calculation of X_{m+1}^R for this case.

For the $Y(4260)$, we first estimate its quasi-two-body partial widths to $Z_c(3900)\pi$, $J/\psi f_0(500)$ and $J/\psi f_0(980)$ based on the analyses of Refs. [22, 23], while the decay width to $\chi_{c0}\omega$ is explicitly given in Ref. [22]. For the $Z_c(3900)\pi$ channel, it accounts 21% of the $J/\psi\pi\pi$ channel, according to Ref. [23]. Regarding the $h_c\pi\pi$ case, the possible important quasi-two-body channel is $h_c f_0(500)$, but then the interaction between h_c and $f_0(500)$ is P -wave, in contrast to the S -wave interaction between J/ψ and f_0 resonances. Therefore we assume that the rest of the decay widths for $Y(4260)$ are saturated by the quasi-two-body decays into $J/\psi f_0(500)$ and $J/\psi f_0(980)$, and then the data points with $m_{\pi\pi} < 0.65$ GeV are identified as $J/\psi f_0(500)$ and the others as $J/\psi f_0(980)$ [23]. From the partial widths we estimate $|\gamma_i|$ and then X_i^R can be calculated and given in Table I

Our results show that the two-body components are dominant for the $D_{s0}^*(2317)$ (a bound state in Ref. [21]), although it also has non-negligible elementariness. This conclusion is consistent with the large N_c trajectory of its pole position [21]. The residues for the $Y(4260)$ are estimated as discussed above. A small value of X^R is found for $Y(4260)$, implying that this state is more like an elementary particle, a conclusion in agreement with Ref. [22] that uses the pole counting rule [29].

The $\Lambda_c(2595)$ was studied in Ref. [30] in the isospin symmetric case. However, its results are unstable under little changes of the isospin limit mass for pions, so that we have made another delicate analysis [31] and the results are stable and shown in Table I. Because of isospin breaking in the threshold energies of the three $\pi\Sigma_c(2450)$ channels we can only calculate compositeness for the lightest one because then condition I is satisfied. Our robust conclusion is that the $\pi^0\Sigma_c^+$ component is small, with the rest of the channels (including $\pi^+\Sigma_c^0$ and $\pi^-\Sigma_c^{++}$) and components playing a dominant role in the resonance composition.

V. CONCLUSIONS

In this work we have derived a probabilistic interpretation of the compositeness relation at the pole of a resonance with a convergent Laurent series for s around Res_P (this is called condition I), with s_P the resonance pole position. It then allows one to calculate the compositeness coefficients for those channels that are open up to $s = \text{Res}_P$. The key point is to transform to a new unitary and symmetric S matrix that shares the same strength in the resonance couplings but has a meaningful compositeness relation. The narrow-resonance case is discussed in detail too. It is shown that the general process can be considered as the analytical extrapolation in s_P from the narrow-resonance case (real s axis) up to the final pole position s_P along a contour parallel to the imaginary s axis.

We have shown that the compositeness relation is an expression of the proper normalization to 1 of a rank 1 projection operator associated to the pole state. It has also been demonstrated in full generality that elementariness for a lower number of channels considered includes the compositeness of other heavier channels. We have furthermore shown that there is indeed an infinity of possible compositeness relations, all them related by an orthogonal-like transformation of the S matrix that is actually driven by a unitary matrix. Of these relations we can select the physically suited one if condition I holds. The narrow resonance case clearly shows that the role of the transformation that allows us to end with such expression is twofold. First, one can isolate in this way a resonant S matrix by removing nonresonant contributions. Second, one can get rid of those phases associated with free particles in the asymptotic states, so as to end with a positive and real X_i .

The final expression for the compositeness X_i is $|\gamma_i^2 \frac{dG(s_P)_i}{ds}|$, being γ_i^2 the residue of the resonance to channel i and $G(s_P)_i$ the unitary two-point loop function for the same channel. As stated, this expression does not hold for all types of resonance poles, unless they satisfy the condition I.

The latter can be stated as the resonance pole s_P must lie in an unphysical Riemann sheet adjacent to the physical one in the region $s_m < s < s_{m+1}$, such that $s_m < \text{Re } s_P < s_{m+1}$, being s_m the threshold of channel m . Then, the previous expression can be applied to calculate X_i for all the channels with $s_i \leq s_m$. If **I** is not met one cannot exclude that X_i is larger than one due to the appearance of extra real positive constants d_i , as it has been discussed.

We have exemplified this method by calculating the two-body components for several resonances of interest, taking their pole position and residues from previous results in the literature. At this point we have to emphasize that we do not pretend to perform an exhaustive study of theoretical analyses/models in the literature for each resonance, but mainly using outputs from serious studies that already provide us with the necessary information to evaluate X_i . In this respect, the uncertainties in X_i are estimated by propagating errors in-

ferred from the literature and added them quadratically, without any attempt to estimate systematic uncertainties that could arise from the explicit models used to extract pole properties. Further applications of this method to other resonances could provide important information.

Acknowledgements We would like to thank E. Oset and T. Hyodo for interesting discussions. This work is supported in part by the MINECO (Spain) and ERDF (European Commission) grant FPA2013-40483-P and the Spanish Excellence Network on Hadronic Physics FIS2014-57026-REDT, the National Natural Science Foundation of China (NSFC) under Grant Nos. 11575052 and 11105038, the Natural Science Foundation of Hebei Province with contract No. A2015205205, the grants from the Education Department of Hebei Province under contract No. YQ2014034, the grants from the Department of Human Resources and Social Security of Hebei Province with contract No. C201400323.

-
- [1] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **115**, 072001 (2015) [arXiv:1507.03414 [hep-ex]].
 - [2] S. Weinberg, Phys. Rev. **130**, 776 (1963).
 - [3] V. Baru, J. Haidenbauer, C. Hanhart, Y. Kalashnikova and A. E. Kudryavtsev, Phys. Lett. B **586**, 53 (2004) [hep-ph/0308129].
 - [4] C. Hanhart, Y. S. Kalashnikova and A. V. Nefediev, Eur. Phys. J. A **47**, 101 (2011) [arXiv:1106.1185 [hep-ph]].
 - [5] T. Hyodo, D. Jido and A. Hosaka, Phys. Rev. C **85**, 015201 (2012) [arXiv:1108.5524 [nucl-th]].
 - [6] F. Aceti and E. Oset, Phys. Rev. D **86**, 014012 (2012) [arXiv:1202.4607 [hep-ph]].
 - [7] T. Sekihara, T. Hyodo and D. Jido, PTEP **6**, 063D04 **2015** [arXiv:1411.2308 [hep-ph]].
 - [8] J. R. Taylor, *Scattering Theory*, John Wiley & Sons, New York, 1972, chap. 6.
 - [9] J. A. Oller and U. G. Meißner, Phys. Lett. B **500**, 263 (2001) [hep-ph/0011146].
 - [10] J. A. Oller and E. Oset, Phys. Rev. D **60**, 074023 (1999) [hep-ph/9809337].
 - [11] L. Alvarez-Ruso, J. A. Oller and J. M. Alarcon, Phys. Rev. D **80**, 054011 (2009) [arXiv:0906.0222 [hep-ph]]; A. Martinez Torres, E. J. Garzon, E. Oset and L. R. Dai, Phys. Rev. D **83**, 116002 (2011) [arXiv:1012.2708 [hep-ph]].
 - [12] Z. H. Guo and J. A. Oller, in preparation.
 - [13] J. A. Oller and E. Oset, Nucl. Phys. A **620** (1997) 438 [Nucl. Phys. A **652** (1997) 407] [hep-ph/9702314].
 - [14] J. A. Oller and M. Albaladejo, Phys. Rev. Lett. **101**, 252002 (2008) [arXiv:0801.4929 [hep-ph]].
 - [15] R. H. Dalitz, *Resonances- models and phenomena. Proceedings of a Workshop held at the Centre for Interdisciplinary Research Bielefeld University, Bielefeld, Germany, April 9-14, 1984*. Edited by S. Alberverio, L. S. Ferreira and L. Streit. Springer-Verlag Berlin Heidelberg 1984. This author also introduced the transformation of Eq. (20) in a narrower context.
 - [16] X. W. Kang, Z. H. Guo and J. A. Oller, arXiv:1603.05546 [hep-ph].
 - [17] K.A. Olive *et al.* (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
 - [18] Z. H. Guo, J. A. Oller and J. Ruiz de Elvira, Phys. Rev. D **86**, 054006 (2012) [arXiv:1206.4163 [hep-ph]].
 - [19] M. F. M. Lutz, S. Leupold, Nucl. Phys. A **813**, 96 (2008) [arXiv:0801.3821 [nucl-th]].
 - [20] Z. H. Guo and J. A. Oller, Phys. Rev. C **87** 3, 035202 (2013) [arXiv:1210.3485 [hep-ph]].
 - [21] Z. H. Guo, U. G. Meißner and D. L. Yao, Phys. Rev. D **92**, no. 9, 094008 (2015) [arXiv:1507.03123 [hep-ph]].
 - [22] L. Y. Dai, M. Shi, G. Y. Tang, H. Q. Zheng, Phys. Rev. D **92**, 014020 (2015) [arXiv:1206.6911 [hep-ph]]. ZHG would like to thank G. Y. Tang for communications on details of this reference.
 - [23] M. Ablikim *et al.* [BESIII Collaboration], Phys. Rev. Lett. **110**, 252001 (2013) [arXiv:1303.5949 [hep-ex]].
 - [24] J. R. Pelaez, Phys. Rev. Lett. **92**, 102001 (2004) [hep-ph/0309292].
 - [25] M. Albaladejo and J. A. Oller, Phys. Rev. D **86**, 034003 (2012) [arXiv:1205.6606 [hep-ph]]. The dependence of the pole position and residue of the $f_0(500)$ as the pion mass varies is given.
 - [26] L. S. Geng, E. Oset, J. R. Pelaez and L. Roca, Eur. Phys. J. A **39**, 81 (2009) [arXiv:0811.1941 [hep-ph]].
 - [27] D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meißner, Nucl. Phys. A **725**, 181 (2003) [nucl-th/0303062].
 - [28] K. Moriya *et al.* [CLAS Collaboration], Phys. Rev. C **87**, no. 3, 035206 (2013) [arXiv:1301.5000 [nucl-ex]].
 - [29] D. Morgan, Nucl. Phys. A **543**, 632 (1992); K. L. Au, D. Morgan and M. R. Pennington, Phys. Rev. D **35**, 1633 (1987).
 - [30] T. Hyodo, Phys. Rev. Lett. **111**, 132002 (2013).
 - [31] Z. H. Guo and J. A. Oller, Phys. Rev. D **93**, no. 5, 054014 (2016) [arXiv:1601.00862 [hep-ph]].